


§1.1 Introduction to Interpolation

§1 Interpolation = "join the dots"

MA378/531 – Numerical Analysis II ("NA2")

January 2017




Suppose that we have a two sets of $n + 1$ real numbers $\{x_i\}_{i=0}^{n+1}$ and $\{y_i\}_{i=0}^{n+1}$, and that the x_i are *strictly* increasing:

$x_0 < x_1 < x_2 < \cdots < x_n$. *Interpolation* problems are of the form:

Find a function p , that is continuous and defined on $[x_0, x_n]$, such that

$$p(x_k) = y_k, \quad \text{for } k = 0, 1, \dots, n.$$

We say that p *interpolates* the points (x_0, y_0) , (x_1, y_1) , ..., (x_n, y_n) .



Why Bother? There are several possibilities, including

- The points belong to an underlying, but unknown function. We wish to establish likely values of f at points other than x_0, x_1, \dots, x_n . The values of f may have been obtained from physical experiments, or numerical procedures (e.g., Newton's method for initial value problems). Or it may be that some values of the function are easily available. For example $2! = 2$, and $3! = 6$, but what about $2\frac{1}{2}!$ or $\pi!$?
- We may know the function, but prefer to work with an interpolant to it. For example, in order to estimate derivatives or integrals of a function.

Applications? In mathematics, from number theory to information theory, and nearly every aspect of numerical analysis. Elsewhere, the methods are used in fields ranging from aircraft design to computer animation.

The main reference for this section is Chapter 6 of Suli and Mayers, See also Lectures 18–20 of Stewart's *Afternotes*.

Finished here 9/1/17

Definition

\mathcal{P}_n is the set of polynomials of degree less than or equal to n and real-valued coefficients, i.e., $p_n \in \mathcal{P}_n$ if

$$p_n(x) = a_0 + a_1x + a_2x^2 + \cdots + a_nx^n,$$

where $a_i \in \mathbb{R}$.

Examples:

The constants $1, -101, \pi$, etc, all belong to \mathcal{P}_0 .

All the polys in \mathcal{P}_1 are of the form $ax + b$
where a, b are real numbers,

$x^3 - x^2 + x + 1, (x-1)(x+1)(x+2)$ belong to \mathcal{P}_3 .

but $\underbrace{(x^2+1)(x^2-1)}_{\in \mathcal{P}_4}$ and $x^3 + x^2 + x + \sqrt{x}$ do not.

Exercise

Find out what a *vector space* is. Convince yourself that \mathcal{P}_n is a vector space.

It is particularly important to note that if p_n and q_n both belong to \mathcal{P}_n , then so too does their sum.

$$\text{Eg } x^2 + 1 \in \mathcal{P}_2 \quad \& \quad x^2 + x \in \mathcal{P}_2 \quad \text{so} \\ 2x^2 + x + 1 \in \mathcal{P}_2.$$

$$\text{Also: } \mathcal{P}_0 \subseteq \mathcal{P}_1 \subseteq \mathcal{P}_2 \subseteq \mathcal{P}_3 \subseteq \dots \subseteq \mathcal{P}_n.$$

$$\text{So, eg } x+1 \in \mathcal{P}_1 \quad \text{and} \quad ax^2 + x + 1 \in \mathcal{P}_2 \\ \text{but } a=0.$$

The Polynomial Interpolation Problem comes in two forms.

The Polynomial Interpolation Problem I (PIP1)

Given is set of points $x_0 < x_1 < \cdots < x_n$, and a set of real numbers y_0, y_1, \dots, y_n , find $p_n \in \mathcal{P}_n$ such that

$$p_n(x_k) = y_k, \quad \text{for } k = 0, 1, \dots, n.$$

The Polynomial Interpolation Problem II (PIP2)

Given is set of points $x_0 < x_1 < \cdots < x_n$, and a function $f : [x_0, x_n] \rightarrow \mathbb{R}$, find $p_n \in \mathcal{P}_n$ such that

$$p_n(x_k) = f(x_k), \quad \text{for } k = 0, 1, \dots, n.$$

Clearly PIP2 is just PIP1 with $y_k = f(x_k)$.

The questions that we must ask (and answer) are

- (i) Is there a solution to the polynomial interpolation problem.
- (ii) Is it unique?
- (iii) How do we find it?
- (iv) How accurate is it? If f is the underlying function (i.e., $f(x_k) = y_k$), can we find an upper bound for

$$\max_{x_0 \leq x \leq x_n} \{|f(x) - p_n(x)|\}?$$

we know that $p_n(x_k) = f(x_k)$, so there is no error at the interpolation points. But what about every other point?

The questions that we must ask (and answer) are

- (i) Is there a solution to the polynomial interpolation problem.
- (ii) Is it unique?
- (iii) How do we find it?
- (iv) How accurate is it? If f is the underlying function (i.e., $f(x_k) = y_k$), can we find an upper bound for

$$\max_{x_0 \leq x \leq x_n} \{|f(x) - p_n(x)|\}?$$

